arXiv:astro-ph/9712305v1 23 Dec 1997

PRIMORDIAL DEUTERIUM ABUNDANCE MEASUREMENTS

Sergei A. LEVSHAKOV
National Astronomical Observatory, Mitaka, Tokyo 181-8558, JAPAN, lev@diamond.mtk.nao.ac.jp
Wilhelm H. KEGEL
Institut für Theoretische Physik der Universität Frankfurt am Main, Postfach 11 19 32, 60054 Frankfurt/Main 11, GERMANY, kegel@astro.uni-frankfurt.de
Fumio TAKAHARA
Department of Earth and Space Science, Faculty of Science, Osaka University, Toyonaka, Osaka 560, JAPAN, takahara@vega.ess.sci.osaka-u.ac.jp

Abstract

Deuterium abundances measured recently from QSO absorption-line systems lie in the range from $\simeq 3 \times 10^{-5}$ to $\simeq 3 \times 10^{-4}$, which shed some questions on standard big bang theory. We show that this discordance may simply be an *artifact* caused by inadequate analysis ignoring spatial correlations in the velocity field in turbulent media. The generalized procedure (accounting for such correlations) is suggested to reconcile the D/H measurements.

An example is presented based on two high-resolution observations of Q 1009+2956 (low D/H) [1,2] and Q 1718+4807 (high D/H) [8,9]. We show that both observations are compatible with D/H $\simeq 4.1 - 4.6 \times 10^{-5}$, and thus support SBBN. The estimated mean value $\langle$ D/H $\rangle \simeq 4.4 \times 10^{-5}$ corresponds to the baryon-to-photon ratio during SBBN $\eta \simeq 4.4 \times 10^{-10}$ which yields the present-day baryon density $\Omega_b h^2 \simeq 0.015$.

1. Introduction

From recent HST observations of the $z_a = 0.7$ absorption-line system toward the quasar Q 1718+4807 Webb *et al.* [8,9] deduced D/H $= 1.8 - 3.1 \times 10^{-4}$. This ratio is significantly higher than that derived from other quasar spectra. For instance, at $z_a = 2.504$ toward Q 1009+2956 the D/H ratio lies in the range from

1.8×10^{-5} to $3.5\,10^{-5}$ [1]. New measurements [2] give a slightly higher range for the D/H value at $z_a = 2.504$ ($3.31 - 4.57 \times 10^{-5}$) which is still incompatible with the results of Webb *et al.*

According to the basic idea of homogeneity and isotropy of the big bang universe the *primordial* deuterium abundance should not vary in space. One can only expect that the D abundance decreases with cosmic time due to conversion of D into ^{3}He and heavier elements in stars. To check whether big bang nucleosynthesis [BBN] has occurred homogeneously or not, precise measurements of absolute values of D/H at high redshift are extremely important. In a series of papers [3-6] we have shown that this task is badly model dependent. Here we present two examples to underline the difficulties of the inverse problem in the analysis of the H+D absorption blends.

2. Results

We consider a cloud of a thickness L with a stochastic velocity field but of homogeneous (H I) density and temperature. The velocity field is characterized by its rms amplitude σ_t and its correlation length $l > 0$. The model is identical to that of [5]. This approach generalizes the standard procedure which assumes no spatial correlations in the velocity field ($l \equiv 0$). – To estimate physical parameters and an appropriate velocity field structure along the line of sight, we used a Reverse Monte Carlo [RMC] technique. The algorithm requires to define a simulation box for 5 physical parameters : N(H I), D/H, T_{kin}, σ_t/v_{th}, and L/l (here v_{th} denotes the thermal width of the hydrogen lines). – The continuous random function of the coordinate $v(s)$ is represented by its sampled values at equal space intervals Δs, i.e. by $\{v_1, v_2, \ldots, v_k\}$, the vector of the velocity components parallel to the line of sight at the spatial points s_j (for more detail, see [5]).

Direct observations of galactic halos at $z > 2$ [7] show that $\sigma_t \simeq 40 \pm 15$ km s^{-1}, if $T_{kin} \simeq 10^4$ K. Our RMC calculations yield for the $z_a = 2.504$ and $z_a = 0.701$ absorption systems $\sigma_t \simeq 26$ km s^{-1} [5,6] which makes the procedure to be adequate, whereas $\sigma_t \simeq 2 - 8$ km s^{-1} found in [1,2] and $\sigma_t \simeq 13$ km s^{-1} found in [8,9] are evidently too low.

To illustrate our results, we show in Fig. 1 H+D Lyα lines observed by the two groups [1,2] and [8,9] and some of the adequate profile fits (χ^2_{min} per degree of freedom $\simeq 1$). The solutions are not unique, however, but depend sensitively on the velocity field configuration along the line of sight (see [5,6], for more examples). Fig. 1 shows that both profiles are compatible with D/H $\simeq 4.1 \times 10^{-5}$ (the uncertainty range is from 4.1×10^{-5} to 4.6×10^{-5} [5,6]). Taking the mean D/H value from this range, one can estimate the baryon-to-photon ratio



Fig. 1. *Upper panel.* An example of a template H+D Lyα profile (dots) at $z_a = 2.504$ toward the quasar Q 1009+2956 representing the normalized intensities and their uncertainties (error bars) in accord with [1]. The results of the RMC minimization are shown by the solid curve (H+D), the dotted curve (H I) and the dashed curve (D I) for the mesoturbulent model with N(H I) $= 2.71 \times 10^{17}$ cm^{-2}, D/H $= 4.11 \times 10^{-5}$, $T_{kin} = 20220$ K, $\sigma_t = 25.8$ km s^{-1}, and $L/l = 4.1$.

Lower panel. HST/GHRS observations [8,9] of the H+D Lyα line at $z_a = 0.701$ toward the quasar Q 1718+4807 (the open circles represent the normalized flux, the spectral resolution is 0.1 Å). Profiles for H I (dotted curve), D I (dashed curve) and H+D (solid curve) are computed by the RMC procedure for N(H I) $= 1.77 \times 10^{17}$ cm^{-2}, D/H $= 4.11 \times 10^{-5}$, $T_{kin} = 15100$ K, $\sigma_t = 26$ km s^{-1}, and $L/l = 3.5$. The χ^2 values (per degree of freedom) of the final fits are shown.

η during SBBN and using the present-day photon density determined from the cosmic microwave background measurements, one can obtain the current baryon density Ω_b : $\eta \simeq 4.4 \times 10^{-10}$ and $\Omega_b h^2 \simeq 0.015$, correspondingly.

We conclude that the reliability of the interpretation of deuterium absorption line observations at high redshift is determined by two factors : (*i*) improvements in the detection equipment, and (*ii*) advances in the theory of line formation in turbulent media.

Acknowledgment – The authors are grateful to John Webb for making available the calibrated HST/GHRS spectra of Q 1718+4807 and acknowledge helpful correspondence and comments by him and Alfred Vidal-Madjar. This work was supported in part by the RFBR grant No. 96-02-16905a.

3. References

1. Burles, S., Tytler, D., 1996, astro-ph 9603069.
2. Burles, S., Tytler, D., 1997, *Astrophys. J.* (submit.), astro-ph 9712109.
3. Levshakov, S. A., Kegel, W. H., 1997, *Monthly Notices Roy. Astron. Soc.* **288**, 787.
4. Levshakov, S. A., Kegel, W. H., Mazets, I. E., 1997, *Monthly Notices Roy. Astron. Soc.* **288**, 802.
5. Levshakov, S. A., Kegel, W. H., Takahara, F., 1997, *Monthly Notices Roy. Astron. Soc.* (submit.), astro-ph 9710122.
6. Levshakov, S. A., Kegel, W. H., Takahara, F., 1997, *Astron. Astrophys.* (submit.) .
7. van Ojik, R., et.al., 1997, *Astron. Astrophys.* **317**, 358.
8. Webb, J. K., et.al., 1997, *Nature* **388**, 250.
9. Webb, J. K., et.al., 1997, astro-ph 9710089.